




## Penaeid shrimp



**Brown shrimp,**  
*Farfantepenaeus aztecus*  
(Ives, 1891)



**Pink shrimp,**  
*Farfantepenaeus duorarum*  
(Burkenroad, 1939)



**White Shrimp,**  
*Litopenaeus setiferus*  
(Linnaeus, 1767)

Three commercially important species of penaeid shrimp occur on both coasts of Florida. The distribution of white shrimp, *Litopenaeus setiferus*, and brown shrimp, *Farfantepenaeus aztecus*, is intermittent in Florida waters. White shrimp do not occur from about St. Lucie Inlet on the Atlantic coast around the southern tip of Florida north to about the mouth of the Ochlockonee River. Brown shrimp do not occur on the gulf coast between Sanibel Island and Apalachicola Bay. All three shrimp species occur in nearshore waters and estuaries and use the estuaries as nursery areas. At various juvenile stages, penaeid shrimp usually inhabit seagrass beds and algal mats within estuaries. Stable isotope studies show young pink shrimp that recruited to the southeastern Gulf of Mexico offshore fisheries are mostly migrants from seagrass meadows (Fry *et al.* 1999). Adult pink shrimp, *F. duorarum*, are most abundant at depths between 35' and 120'. White shrimp are most abundant in waters shallower than 90', and brown shrimp are most abundant in waters less than 180'. White shrimp are typically distributed in areas of low salinity over organic-rich, mud bottoms. Brown shrimp are found on similar bottoms but in higher salinities. Pink shrimp occur on more coarse sediments and in a wide variety of salinities (Steele unpublished ms.). White shrimp grow rapidly until about 6.3 inches total length (TL). Peak growth rates are 0.8 inches/month during summer. Brown shrimp can grow at peak rates of 1.8 inches/month during spring; pink shrimp peak growth rates have been reported to exceed 2.0 inches/month.

All three species mature during their first year. Sizes at maturity are about 5.5 inches TL for white and brown shrimp and about 3.3 inches TL for pink shrimp. Spawning occurs in relatively deep water for brown shrimp (49'–360') and pink shrimp (13'–160'), and in nearshore waters (20'–80') for white shrimp. White shrimp spawn during April–October. Pink and brown shrimp can spawn year-round, especially in deeper or more southern waters. Peak spawning occurs during February and March for brown shrimp and during spring, summer, and fall for pink shrimp.

Reported commercial penaeid shrimp landings totaled 11,809,484 pounds in Florida during 2007. Most landings (56%) were made on the gulf coast. The geographic distribution of landings for each species was different. The greatest landings of brown shrimp were reported in the Florida Panhandle region and in Hillsborough and Lee Counties on the gulf coast, and in Nassau, Duval, and Brevard Counties on the Atlantic coast (Fig. 1a). Pink shrimp were landed mostly in Dade, Monroe, Lee, Pinellas, Hillsborough, Gulf, and Franklin Counties (Fig. 1b). White shrimp were landed mostly in northeast, east-central Florida (Nassau-Brevard Counties), and to a lesser extent in Bay and Franklin Counties on the gulf coast (Fig. 1c) in areas adjacent to extensive saltwater marshes and high freshwater run-off.

The 2007 total landings of penaeid shrimp were 35% lower than the average landings in the previous five years (2002–2006) and were 41% lower than the 1986–2007 historical average landings (Fig. 2). Commercial landings of penaeid shrimp increased on the Atlantic coast from a low of about 2.5 million pounds in 1984 to a peak of about 5.4 million pounds in 1999. Recent commercial landings figures indicate that there were 4.8 million pounds landed in 2002, 5.3 million pounds in 2004, 4.7 million pounds in 2005, and about 5 million pounds in both 2006 and 2007 (Fig. 2). Gulf coast landings increased dramatically, from about 10.6 million pounds in 1992 to 24.6 million pounds in 1996, and then dropped to 17.8 million pounds in 1997 before rebounding to 22.3 million pounds in 1998. Gulf landings declined after 1998, averaged 13.4 million pounds during the period 1999–2005, and declined to about 7 million pounds in 2007. Competition from foreign imports and increased fuel and operating costs contributed significantly to the declines in penaeid shrimp landings by commercial shrimpers over the last 9 years.

Standardized annual landings rates for all three species generally increased or held steady on both the Atlantic and gulf coasts from 1999–2006, but landings rates have declined on the gulf coast during 2007 for all three species and for pink shrimp on the Atlantic coast [Fig. 3(a)-(f)]. Brown shrimp landings rates have fluctuated with an increasing trend on both coasts from 1992–2007 [Fig. 3(a)-(b)]. Landings rates for pink shrimp fluctuated on both coasts, but peaked in the mid 1990s, declined then until 2000 and exhibited an increasing trend during 2003–2006, and have declined back to 2005 levels [Fig 3 (c)-(d)]. White shrimp landings rates on the Atlantic coast fluctuated without a trend until 2003, and have since increased [Fig. 3(e)]. On the gulf coast, white shrimp catch rates increased from 1992 to 1997, declined between 1999 and 2001 increased significantly during 2003–2006, and declined significantly in 2007 [Fig. 3(f)].

The index of relative abundance of young-of-the-year (YOY) pink shrimp fluctuated without trend on the Atlantic coast from 1996 through 2004 after which abundances declined through 2007 (Fig. 4a). Recruitment of gulf coast YOY pink shrimp shows a discrete declining trend since from 2000 to 2005, the abundance of YOY pink shrimp appears to be rebounding in 2006 and 2007 (Fig. 4b).

An assessment of the condition of U.S. gulf and South Atlantic penaeid shrimp stocks suggests that they are all harvested at or slightly in excess the fishing mortality rates associated with maximum yield-per-recruit (Steele unpubl. data). Increasing the size-at-entry to the fishery could increase the yield and value of the landings for all three shrimp species. Available data do not suggest a strong link between parent stock abundance and subsequent abundance of their progeny. Regardless, estimated spawning potential ratios were estimated to be 4%–12% for brown shrimp and 13%–39% for white shrimp during 1970–1987 (Nance *et al.* 1989).

Nance (1999) found that the parent stock levels for brown shrimp in the Gulf of Mexico were up in 1998 at over 300 million age-7<sup>+</sup>-month-old shrimp for November through February, well above the 125 million overfishing threshold and the highest level since 1994. For white shrimp, the parent stock number had been highly variable since the mid 1980s and the number dropped slightly in 1998 to around 800-million individuals age-7<sup>+</sup> months for May through August. However, this level is still well above the 330 million individuals overfishing threshold. Pink shrimp parent stock numbers were up in 1999 following a slight decline in 1998 to nearly 250-million age-5<sup>+</sup> -month-old individuals for July through June and were well above the 100 million overfishing threshold.

Both regional federal councils, the South Atlantic Fishery Management Council and the Gulf of Mexico Fishery Management Council, have established fishery management plans for

shrimp (GMFMC 1981; SAFMC 1993). The main objectives of these plans were to delay harvest of shrimps through season and area closures, reduce bycatch, and minimize gear conflicts. The SAMFC's shrimp fishery management plan was instituted to protect the white shrimp stock from over-harvest after severe winter cold-kills. This plan allows for the closure of the Exclusive Economic Zone after severe winter kills and requires permits as a first step toward possible limited entry.




Figure 1 (a)-(c). Geographic distribution of penaeid shrimp landed commercially during 2007 by species and county.




Figure 2. Total annual commercial landings (pounds) of penaeid shrimp on the Atlantic and gulf coasts of Florida, 1982–2007.




Figure 3 (a)-(f). Annual standardized catch rates for penaeid shrimp in Florida. Commercial landings rates (pounds/trip) 1992-2007: (a) Atlantic Coast brown shrimp; (b) Gulf Coast brown shrimp; (c) Atlantic Coast pink shrimp; (d) Gulf Coast pink shrimp; (e) Atlantic Coast white shrimp; (f) Gulf Coast white shrimp.





Figure 4(a)-(b). Proportion of fishery-independent-monitoring sets that captured pink shrimp from 1996-2007. Young-of-the-year (YOY): (a) Atlantic coast; (b) Gulf coast.